

Coupled Harmonic Oscillators

Applied Statistical thermodynamics

November 10, 2015

1 Conclusions [1]

1.1 Question 3

From our simulation experiment in question 3, it follows that the probability distribution for a given state k of a single harmonic oscillator reads:

$$P(k) = \frac{\exp[-\beta E_k]}{q} \quad (1)$$

with constants β and q . In statistical thermodynamics the numerator $\exp[-\beta E_k]$ is called the Boltzmann factor. The constant q is called the molecular partition function or molecular partition sum, defined as the sum of Boltzmann factors over all states of the oscillator.

$$q = \sum_{k=0}^{\infty} \exp[-\beta E_k] \quad (2)$$

Substituting $x = \exp(-\beta \hbar \omega)$ we can rewrite the partition sum of q of a harmonic oscillator as a geometric series employing:

$$\sum_{i=0}^{\infty} x^i = \frac{1}{1-x} \quad \text{for } |x| < 1 \quad (3)$$

thus:

$$q = \frac{1}{1 - \exp[-\beta E_k]} \quad (4)$$

The average energy $\langle E_i \rangle$ of a single oscillator can be related to β using the same substitution as in eq. (3) and employing :

$$\sum_{i=0}^{\infty} i x^i = \frac{x}{(1-x)^2} \quad \text{for } |x| < 1 \quad (5)$$

which is eq (3) differentiated with respect to x and multiplied by x , leads to the following relation:

$$\langle E_i \rangle = \sum_{k=0}^{\infty} \hbar \omega k P(k) = \hbar \omega \frac{\sum_{k=0}^{\infty} k \exp[-\beta \hbar \omega k]}{\sum_{k=0}^{\infty} \exp[-\beta \hbar \omega k]} = \frac{\hbar \omega}{\exp[-\beta \hbar \omega] - 1} \quad (6)$$

The latter means that for big enough number of oscillators, a single oscillator i is surrounded by a "heath bath". The temperature follows from the total energy E , which in turn determines $\langle E_i \rangle$ and β . Thus, from the average energy $\langle E_i \rangle$ and the relation $\beta = 1/k_B T$ we can estimate the temperature of the simulated system.

1. For a number of system energies (E_t) use simulation data to find $\langle E_i \rangle$ and β . Check your results with eq. (6)

1.2 Question 4

From our simulation experiment in question 4, it follows that the entropy increase in time and reaches maximum value when we start from the less likely distribution.

References

- [1] Thijs J.H. Vlugt, Jan P.J.M. van der Eerden, Marjolein Dijkstra, Berend Smit, and Daan Frenkel. *Introduction to Molecular Simulation and Statistical Thermodynamics*. Delft, The Netherlands, 2008.